



BEEF HEIFER PROTOCOLS - 2011

HEAT DETECTION


1 Shot PG



7-day CIDR®-PG




MGA®-PG



HEAT DETECT & TIME AI (TAI)


Select Synch + CIDR® & TAI

Heat detect and AI day 7 to 10 and TAI all non-responders 72 - 84 hr after PG with GnRH at TAI.




MGA®-PG & TAI

Heat detect and AI day 33 to 36 and TAI all non-responders 72 - 84 hrs after PG with GnRH at TAI.



14-day CIDR®-PG & TAI


Heat detect and AI day 30 to 33 and TAI all non-responders 72 hrs after PG with GnRH at TAI.



FIXED-TIME AI (TAI)*


7-day CO-Synch + CIDR®

Perform TAI at 54 ± 2 hr after PG with GnRH at TAI.




MGA®-PG

Perform TAI at 72 ± 2 hr after PG with GnRH at TAI.



14-day CIDR®-PG

Perform TAI at 66 ± 2 hr after PG with GnRH at TAI.



COMPARISON OF PROTOCOLS FOR BEEF HEIFERS

HEAT DETECTION	COST	LABOR
1 Shot PG	Low	High
7-day CIDR®-PG	High	Medium
MGA®-PG	Low	Low/Medium

HEAT DETECT & TAI

Select Synch + CIDR® (TAI non-responders 72-84 hr after PG)	High	Medium
MGA®-PG (TAI non-responders 72-84 hr after PG)	Medium	Medium
14-day CIDR®-PG (TAI non-responders 70-74 hr after PG)	High	Medium

FIXED-TIME AI (TAI)

CO-Synch + CIDR® (TAI 54 ± 2 hr after PG with GnRH at TAI)	High	Medium
MGA®-PG (TAI 72 ± 2 hr after PG with GnRH at TAI)	Medium	Medium
14-day CIDR®-PG (TAI 66 ± 2 hr after PG with GnRH at TAI)	High	Medium

* The times listed for "Fixed-time AI" should be considered as the approximate average time of insemination. This should be based on the number of heifers to inseminate, labor, and facilities.

GnRH Cystorelin®, Factrel®, Fertagyl®, OvaCyst®

PG estroPLAN®, Estrumate®, In-Synch®, Lutalyse®, ProstaMate®